

THE SPEED OF GRAVITY: WHAT A THEORY SAYS

Antonio Alfonso-Faus
E.U.I.T. Aeronáutica, Spain

Following a quantum-gravity approach we use a gravitational quantum defined elsewhere as well as an effective gravitational “cross section” in conjunction with Mach’s Principle and the de Broglie wavelength concept. We find the speed of gravity v_g related to a gravitational mass m as given by $v_g \approx c(M_u/m)^{1/2}$, where c is the speed of light and M_u the mass of the seeable Universe. For the solar system we get $v_g \approx 10^{11} c$, which agrees very well with experimental results. For the Universe $v_g \approx c$.

Keywords: Gravitation; Speed; Relativity; Faster-than-light; Tachyons; Quantum Gravity.

1 Introduction

The concept of “action at a distance” for gravity implies an infinite propagation speed. We believe that there are no infinities of any kind in Nature: there is always a natural way to avoid such infinities. Many arguments can be given in favor of a finite, but much larger than the speed of light c , speed of gravity [1] including experimental results. If gravity is due to some kind of interaction (as a quantum-gravity approach would demand) then the speed of propagation of this interaction should be determined by a theory of gravity quanta [2]. We know that there is no concept of force for gravity under a general relativity (GR) approach. But GR is not a quantum theory. If one includes quanta for gravity then the propagation of the interaction implying a force (a change of momentum with time by an absorption-emission process) seems to be the appropriated approach. And then the question of the speed of this propagation arises and a value much faster than light seems to be the case [1] on an experimental basis.

Working with a gravity quantum and a gravitational cross section [2] we will find here the speed of gravity using Mach’s Principle and the de Broglie wavelength concept. It turns out that the speed of gravity in general is many orders of magnitude much larger than the speed of light (except for the Universe itself). To all intents and purposes this is practically an infinite value. This result is in contrast with a

relativity principle that considers the speed of light as an absolute maximum. But relativity is consistent with faster-than-light particles (tachyons). This means that our theory predicts a propagation of gravitational interactions through tachyons in the form of gravity quanta.

2 The Speed of Gravity

To determine the speed of gravity from a gravitational mass m one needs the concept of a quantum of gravity of mass m_g [2]:

$$m_g = \frac{\hbar}{c^2 t} \quad (1)$$

and a gravitational cross section σ_g [2]:

$$\sigma_g = \frac{Gm}{c^2} \cdot ct \quad (2)$$

This gravitational cross section is the effective area of gravitational interaction for the mass m . It is given by the product of its gravitational radius Gm/c^2 and the size of the seeable Universe ct (t the age of the Universe). Interpreting Mach's Principle under the view that the rest energy of any mass mc^2 is due to its gravitational potential with respect to the mass of the rest of the seeable Universe M_u :

$$\frac{GM_u m}{ct} = mc^2 \quad (3)$$

one has

$$\frac{GM_u}{c^2} = ct \quad (4)$$

Then the gravitational cross section of any mass given in (2) is the product of two gravitational radii Gm/c^2 and GM_u/c^2 .

The uncertainty principle for a quantum of gravity of mass m_g and speed v_g gives:

$$m_g v_g (\sigma_g)^{1/2} \approx \hbar \quad (5)$$

This is as much as to say that the de Broglie wavelength for the quantum of gravity has about the size of the effective gravitational cross section:

$$(\sigma_g)^{1/2} \approx \frac{\hbar}{m_g v_g} \quad (6)$$

Then from (1), (2) and (5) we get

$$v_g^2 \approx \frac{c^5}{G} \cdot \frac{t}{m} \quad (7)$$

and using Mach's Principle in the form in (4) one finally has:

$$v_g \approx c \sqrt{\frac{M_u}{m}} \quad (8)$$

This is the speed of gravity as related to any mass m . For the Universe as a whole we have $m = M_u$ and therefore $v_g \approx c$. There are about 10^{11} galaxies in the seeable Universe. Then gravity from a galaxy has a speed of about $3 \cdot 10^5 c$. There are about 10^{11} equivalent solar systems in a galaxy. Then gravity inside a solar system has a speed of about $10^{11} c$. Now, Van Flandern in [1] finds from experiments a lower limit for the value of the speed of gravity $v_g \geq 2 \cdot 10^{10} c$, which implies a complete agreement between our theory and the experiments.

3 Conclusions

Under a gravity quanta approach the speed of gravity v_g is determined to be much faster than light, $v_g \approx c(M_u/m)^{1/2}$. Then gravity propagates as tachyons, consistent with the relativity principle. The experimental results do not contradict this theoretical deduction.

References

- [1] T. Van Flandern, *Physics Letters A* **250** (1998) 1-11, Elsevier Science.
- [2] A. Alfonso-Faus, *Physics Essays*, Vol. **12**, N. 4, (1999) 673-677.